

Remember: take your blue inhaler **before** you come into contact with any of your triggers if needed and regularly in response to symptoms if you have a cold.

My Triggers are:

-
-
-
-
-
-
-
-

Common Triggers are:

- Viruses
- Changes in weather
- House dust mites
- Animal fur, feathers and their bedding
- Foods
- Exercise
- Upset, distress, and emotions
- Smoke – cigarettes and fires

Additional Comments:

.....

.....

.....

.....

.....

.....

.....

**Your Asthma Nurse's name
and telephone number is:**

.....

.....

**Your doctor's name
and telephone number is:**

.....


.....



**Asthma Management
Plan For**

Best Peak Flow

Date



Recommended websites
www.beatasthma.co.uk

Asthma+LungUK at:
www.asthma.org.uk

<https://uk-air.defra.gov.uk/forecasting/>

This leaflet is intended for colour printing.

Please take this with you when you visit your
doctor or asthma nurse.

Green zone - Good



Your asthma is under control if:

- your breathing feels good
- you have no cough or wheeze
- your sleeping is not disturbed by coughing
- you are able to do your usual activities
- you are not missing school
- if you check your Peak Flow, it is around your best

BEST PEAK FLOW

Green Zone Action - take your normal medications

Your preventer inhaler is a colour and is called

You take puffs/sucks every morning and every night even when you are well.

Other asthma medications you take are:

.....
.....
.....

Your reliever inhaler is a colour and is called

You take puffs/sucks up to 3 times in a week for symptoms and before exposure to your triggers (see your list) if needed.

If you are needing to use your reliever inhaler more than 3 times per week for symptoms

Move to the AMBER ZONE

Amber zone - Warning



If you are using your blue inhaler more than 3 times per week for symptoms or you often wake at night with a cough or wheeze, arrange a review with your asthma nurse or GP.

Warning signs that your asthma is getting worse:

- you have symptoms (cough, wheeze, 'tight chest' or feel out of breath)
- you need your reliever inhaler more than usual
- your reliever is not lasting **four hours**
- your peak flow is down by a third

PEAK FLOW 1/3 DOWN

Amber Zone Action - continue your normal medicines AND

- Take **2 puffs** of the BLUE inhaler with your spacer 1 puff at a time. Keep doing this every 10 minutes if you still have symptoms up to a total of 6 puffs
- You can do this every 4 hours but **must** make an appointment at your GP surgery within the next 24hrs even if you feel better.
- If you need to do this more than every 4hrs, you must see your GP today or go to A&E
- Start keeping a record of your symptoms and peak flow readings to take to the Doctor

IMPORTANT:

- If after your **6 puffs** you still have increasing wheeze or chest tightness

Move to the RED ZONE

Red zone - Severe



- you are still breathing hard and fast
- you still feel tight and wheezy
- you are too breathless to talk in a sentence
- you are feeling frightened and exhausted

Other serious symptoms are:

- colour changes - very pale / grey / blue
- using rib and neck muscles to breath, nose flaring

Red Zone Action

Take 10 puffs of the blue inhaler via a spacer and call 999

- Asthma can be life threatening
- Do not attempt to do a peak flow
- Whilst waiting for the ambulance and using your spacer, take 1 puff at a time of your blue inhaler, breathing at a normal rate for 4-5 breaths, every 30 seconds.
- Stay where you are and keep calm
- If your child becomes unresponsive and has an adrenaline pen for allergies-use it now.

Additional comments or information

.....
.....
.....
.....
.....
.....
.....